

2016-09

Straylight from explanted intraocular lenses

Labuz, G

<http://hdl.handle.net/10026.1/12547>

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

OPEN ACCESS

ARVO Annual Meeting Abstract | September 2016

Straylight from explanted intraocular lenses

Grzegorz Labuz; ELENI PAPADATOU; Thomas J T P Van Den Berg; Fernando Vargas-Martin; Norberto Lopez-Gil; Nicolaas J Reus

— Author Affiliations & Notes

Grzegorz Labuz

Rotterdam Ophthalmic Institute, Rotterdam, Netherlands

ELENI PAPADATOU

University of Valencia, Valencia, Spain

Thomas J T P Van Den Berg

Royal Netherlands Academy of Arts and Sciences, Amsterdam, Netherlands

Fernando Vargas-Martin

Facultad de Óptica y Optometría, Universidad de Murcia, Murcia, Spain

Norberto Lopez-Gil

Facultad de Óptica y Optometría, Universidad de Murcia, Murcia, Spain

Nicolaas J Reus

Department of Ophthalmology, Amphia Hospital, Breda, Netherlands

Footnotes

Commercial Relationships **Grzegorz Labuz**, None; **ELENI PAPADATOU**, None; **Thomas Van Den Berg**, Royal Netherlands Academy of Arts and Sciences, Amsterdam, the Netherlands (P); **Fernando Vargas-Martin**, None; **Norberto Lopez-Gil**, None; **Nicolaas Reus**, None

Support The AGEYE project (608049), the Marie Curie Initial Training Network program (FP7-PEOPLE-2013-ITN), granted by the European Commission, Brussels, Belgium.

Investigative Ophthalmology & Visual Science September 2016, Vol.57, No Pagination Specified. doi:

Abstract

Purpose : Several studies found that visual straylight does as a rule not return to the level normal for young eyes after cataract extraction and IOL implantation (J Cataract Refract Surg. 41(7):1406-15). Ten percent of pseudophakic patients suffer from disabling straylight elevation. The reason of the observed increase is unknown.

Although it has been reported that scattering of a new intraocular lens (IOL) is low, this may change once the IOL is implanted in the eye. In the present study light scattering expressed in term of straylight of explanted IOLs was evaluated.

Methods : A total of 6 explanted IOLs were studied using a commercial straylight meter (C-Quant, Oculus) adapted for in-vitro analysis of scattering from IOLs (Biomed Opt Express. 19;6(11):4457-64). The adaptation involved exposure of the IOL to the C-Quant target, and observation of the test field by experienced observers. The eyes of the observers were not exposed to the straylight source of the C-Quant, so that only straylight caused by the IOL was evaluated (the eye acting as a detector only). To study whether large or small (compared to wavelength) particles predominate in light scattering of the explanted IOLs, interference filters (IF) were used of 468, 550, and 650 nm. This allows to analyze the particles' size by means of straylight-wavelength dependency. Because of the narrow bandwidth of the IFs (10 nm) a camera was used as light amplifier to enable the suprathreshold psychophysical test, as intended with the C-Quant. Additionally, the camera evaluation was done without the IFs (white light) and compared with results without camera use.

Results : A good correspondence between the straylight measurements with and without camera use was found ($R^2=0.98$). The mean \pm SD (range) straylight of the explanted IOLs was $0.37 \pm 0.65 \log(s)$ (-0.35 to 1.09 log(s)). As 3 of the analyzed IOLs showed very low straylight levels, they were excluded from spectral analysis. The spectral analysis of the other 3 IOLs showed rather weak spectral effects, much less than Rayleigh type scattering.

Conclusions : Although most of the explanted IOLs preserve their low scattering properties, some of the analyzed lenses showed increased straylight. This may result in elevated sensitivity to glare, and consequently lead to patient dissatisfaction. Particles larger than wavelength appear to dominate in light scattering from the explanted IOLs. The origin of these particles must be studied.

This is an abstract that was submitted for the 2016 ARVO Annual Meeting, held in Seattle, Wash., May 1-5, 2016.

This work is licensed under a [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License](https://creativecommons.org/licenses/by-nc-nd/4.0/).

